arXiv:gr-qc/9805090v1 26 May 1998

*

Classification of Inflationary Potentials

Mikel Susperregi
Astronomy Unit, School of Mathematical Sciences, Queen Mary & Westfield College, University of London, London E1 4NS, United Kingdom
(February 7, 2008)

Brans-Dicke gravity is remarkable not only in that General Relativity and Mach's Principle find a common enlarged scenario where they are mutually consistent, but also in that it provides a very interesting quantum cosmological model within the inflationary paradigm. The interplay between the Brans-Dicke scalar Φ and the inflaton field σ plays an important rôle during the course of inflation, and although the dynamics as such is governed by the potential, the onset and the end of inflation are determined by the values of both fields jointly. The relative position of the beginning– and end-of-inflation curves (BoI and EoI respectively) is the most relevant factor in determining the resulting quantum cosmological scenario. The classification of potentials that is given in this paper is based on the criterion of whether the BoI and EoI boundaries enclose a finite or infinite area in the (σ,Φ) plane where inflation takes place. It is shown that this qualitative classification distinguishes two classes of potentials that yield very different cosmologies and it is argued that only those theories in which BoI and EoI enclose a finite area in the (σ,Φ) plane are compatible with our observable universe.

PACS: 98.80.Cq gr-qc/9805090

One of the key features that makes the inflationary paradigm successful is that it provides a mechanism to explain the origin of fluctuations and predict a scale-invariant Harrison-Zel'dovich spectrum at the end of inflation [1,2]. Chaotic inflation [2,3] in particular describes the universe as a statistical ensemble of regions, thus enabling us to envisage a quantum cosmological scenario whose properties are entirely dependent on the potential of the scalar field. In these theories, homogeneous regions in an ensemble of universes are subdivided into further regions where the scalar field σ takes a large number of values. The scalar field evolves according to a stochastic equation $\dot{\sigma} = \dot{\sigma}_S + \xi$ that describes its Brownian motion, as formulated in [4]. The value of the field is hence the combination of the slow-roll solution σ_S, that is the solution of $\dot{\sigma}_S = -(3H)^{-1}V'$ and is due to the contribution of the fields over scales $\gtrsim H^{-1}$, and quantum fluctuations within $\lesssim H^{-1}$ that are related to the stochastic term ξ. The Starobinsky noise ξ follows a Gaussian distribution centered around the classical solution σ_S and its variance is $\sim H/2\pi$.

An arbitrary region ϵH^{-1} of an initial homogeneous volume $\sigma = \sigma_A$ that undergoes a quantum jump $\delta\sigma$ in a timestep δt is taken to a new classical trajectory that corresponds to the hypersurface $\sigma = \sigma_A + \delta\sigma$, and the forthcoming fluctuations at the next timestep within this region are gaussianly distributed around the new classical value. The distribution of the so-called *coarse-grained* field is then governed by the Fokker-Planck equation and it is only dependent on the form of the potential [4–6]. The inflationary expansion of each region that results from this continuous subdivision of homogeneous regions due to quantum jumps depends on the local value of σ, and thus quantum fluctuations that take the scalar field to larger values expand much faster than those that take it to smaller values. After a sufficiently long span of inflation, quantum fluctuations take the scalar field step by step to larger and larger values, and the volume of the universe becomes dominated by regions with values of σ close to the largest permissible for a given potential. This situation is summarized in the simplest terms in Fig. 1. The starting point in this example is a homogeneous bubble $\sigma = \sigma_0$, shown at the top of the figure. After a brief lapse of inflation δt, quantum fluctuations divide the region in two halves; these are homogeneous regions resulting from a positive quantum fluctuation (right, denoted $\uparrow$), and a negative one (left, $\downarrow$) around $\sigma = \sigma_0$. Both fluctuations are for simplicity of the same amplitude $\delta\sigma$, as will all subsequent fluctuations. The volume of the region $\uparrow$ on the right expands faster than the region $\downarrow$ on the left and is larger by a factor $\sim e^{6H'\delta\sigma\delta t}$. After a second lapse of time δt, the two regions $\uparrow$ and $\downarrow$ undergo further divisions in very much the same way and yield four homogeneous regions from the subdivision of the previous two in positive and negative quantum fluctuations of the same amplitude. These volumes are labelled $\downarrow\downarrow$,$\downarrow\uparrow$, $\uparrow\downarrow$,$\uparrow\uparrow$. The relative sizes of the bubbles in the figure are not to scale but are only intended to illustrate that some volumes are bigger than others (typically by many orders of magnitude). Certainly, in this representation where we

have chosen identical fluctuation amplitudes $\delta\sigma$, the regions $\uparrow\downarrow$ and $\downarrow\uparrow$ are of equal size, so their volumes add up. A more realistic situation of the phenomenon depicted in the toy representation of Fig. 1 is that the quantum fluctuations $\delta\sigma$ follow a Gaussian distribution, and therefore the many resulting regions of the type $\uparrow\downarrow$, $\uparrow\downarrow\uparrow\downarrow$ are really part of vast number of regions for which positive and negative fluctuations even out and therefore lay on the trajectory that follows the classical evolution of $\sigma = \sigma_0$.




FIG. 1. Simple sketch of the division of homogeneous regions due to quantum jumps. Positive quantum jumps ($\uparrow$) are located to the right of each subdivision, whereas negative jumps ($\downarrow$) of the same magnitude are located to the left. The total volume is dominated at every stage by the regions with largest values of the fields.




FIG. 2. Powerlaw potential. Beginning- (BoI) and end-of-inflation (EoI) boundaries are represented by the slashed and dotted straight lines respectively. BoI is given by $V(\sigma) \approx M_P^4$ whereas EoI is determined by $\dot{\sigma}^2 \approx 2V(\sigma)$. The configuration at A is sustained by quantum jumps from smaller values of σ, such as B.

As was shown in [7], following the argument sketched above, the largest volume of the universe is occupied by values of σ that correspond to the maximum values of the potential, i.e. in the neighbourhood of the Planck or beginning-of-inflation boundary (BoI), $V(\sigma) \approx M_P^4$. It can be seen in Fig. 2, for the case of a powerlaw potential, that the Planck boundary sets a delimiter to the rate of expansion that corresponds to a certain $\sigma \approx \sigma_{\max}$ and the volume of the universe is dominated by regions where the scalar field takes this value. Whereas the field will classically slow-roll down the potential, quantum jumps that take σ back up to larger values will be enhanced as they create volumes that grow faster, and therefore, as is shown in Fig. 2, the field that evolves classically from the initial location A to B is typically taken back up to A via a quantum transition. Quantum fluctuations retain the field in the neighbourhood of BoI for as long as possible, and therefore an arbitrary region of the universe that has undergone thermalization is most likely to result from a region where the scalar field has stayed in the neighbourhood of BoI for a long time and then rolled down the potential as quickly as possible, via coherent quantum jumps, to cross the end-of-inflation (EoI) boundary.

In this simplest case of inflation driven by one scalar field σ, the BoI is a scale inherent to the physics that depends on the value of M_P, which is a constant of Nature, and the amplitude of V is tuned to predict the right order of magnitude in the spectrum of fluctuations, etc. On the other hand, scalar-tensor theories of gravity (see e.g. [8] for a review) provide a means to investigate M_P in terms of a scalar field, e.g. the Brans-Dicke (BD) field Φ, so that M_P varies from one region to another in the universe like a coarse-grained field, in much the same way as the inflaton in the chaotic model; the combination of these G-varying theories with chaotic inflation yields interesting scenarios where the interplay between Φ and the inflaton σ determines the distribution of M_P in an ensemble of universes in terms of the potential of the scalar field, and therefore, ultimately in the particle physics involved. Two interesting scenarios that result from the combination of BD gravity and chaotic inflation are induced [9] and extended [10] inflation. The difference between both is that in the former the BD field has a non-vanishing potential, whereas the latter has not. For simplicity in this paper we will investigate extended inflation only, and the same analysis applies to induced inflation in the straightforward way.

The extended inflation action is [10]

$$S = \int d^4x\,\sqrt{-g}\left[\Phi R - \frac{\omega}{\Phi}(\partial\Phi)^2 - \frac{1}{2}(\partial\sigma)^2 - V(\sigma)\right], \quad (1)$$

where $V(\sigma)$ is the inflation potential, Φ is the BD field, ω is the BD coupling, that we take to be constant. The slow-roll variational equations in an FRW background take the form

$$\frac{\dot{\Phi}}{\Phi} = 2\frac{H}{\omega}\,, \quad (2)$$

$$\dot{\sigma} = -\frac{1}{3H}V'(\sigma)\,, \quad (3)$$

$$H^2 = \frac{1}{6\Phi}V. \quad (4)$$

The BoI boundary is $V(\sigma) = M_P^4(\Phi)$ (where $M_P^2(\Phi) \equiv 16\pi\Phi$), and EoI is given by

$$\frac{1}{2}\dot{\sigma}^2 + \omega\frac{\dot{\Phi}^2}{\Phi} \approx V(\sigma). \tag{5}$$

With the aid of (2) EoI is rewritten as

$$\Phi = \left(\frac{3\omega - 2}{\omega}\right)\left(\frac{V}{V'}\right)^2. \tag{6}$$

From (2) the following conservation law follows [12,13]

$$\frac{d}{dt}\left[\omega\Phi + \int d\sigma \frac{V(\sigma)}{V'(\sigma)}\right] = 0, \tag{7}$$

which emerges due to the simplicity of the slow-roll equations and does not correspond to a symmetry inherent in the physics.

The functional form of BoI and EoI suggests the following classification of V in two classes, based on the relative position of the boundaries of inflation:

- **Class I** *The area of inflating regions enclosed between the boundaries BoI and EoI is finite.* In this case, the BoI and EoI curves intersect at non-trivial values of the fields.

- **Class II** *The area enclosed by BoI and EoI is infinite.*

In the following we illustrate these classes with some examples, and argue that this qualitative classification distinguishes a very important feature that leads to two entirely different cosmological scenarios.

Firstly we investigate powerlaw potentials $V(\sigma) = \lambda/(2n)\,\sigma^{2n}$ of both *class I* and *class II*. Fig. 3 shows the BoI and EoI boundaries for a powerlaw potential for several n. The EoI boundary is a parabola for all n (and only its amplitude is scaled by a factor $1/(4n^2)$). The BoI boundary on the other hand strongly depends on n, i.e. $\Phi \sim \sigma^n$. For $n = 1, 2$ BoI does not intersect EoI, and both boundaries span an infinite region on the (σ,Φ) plane. Therefore $n = 1, 2$ are *class II* potentials, whereas $n > 2$ are on the other hand *class I*. The classical trajectories (7) are inverted parabolas and the fields move in the direction $A \to B$, in concentric parabolas, along the segment of these curves that is contained within BoI and EoI. Initially, the fields that start out at (σ_0,Φ_0) move along the parabola that crosses this point, though quantum fluctuations soon disperse the motion of the fields to concentric trajectories. In turn, regions that have switched to other classical trajectories via quantum jumps will predominantly follow the slow-roll trajectory, and they are further subdivided in other domains due to later quantum fluctuations that take the fields to other adjacent classical trajectories. The end result, qualitatively, is that regions that move to larger and larger values of σ dominate the total volume of the universe. The important point to note here, as opposed to the σ-only scenario sketched in Fig. 1 where BoI is a fixed quantum scale, is that potentials of the second class, such as $n = 1, 2$, allow σ to grow indefinitely while remaining in the region enclosed by BoI and EoI. In this situation, any arbitrarily large value of the fields is much likelier and predominant than smaller values, but at the same time its likelihood is negligible in comparison to larger values. Hence, in the scenario described by potentials of *class II*, the fields have no typical values and the dominant contribution to the total volume is one where both fields blow up.*




FIG. 3. BoI and EoI boundaries for powerlaw potentials. EoI is thick solid line, scaled to the same amplitude in all three cases $n = 1, 2$ and $n > 2$. BoI is given by: slashed line ($n = 1$); solid line ($n = 2$); dotted line ($n > 2$). Classical trajectories are parabolae concentric to AB. BoI and EoI intersect in the case of $n > 2$.

In contrast to this scenario, potentials $n > 2$ describe a very different universe. As it can be seen in Fig. 3, $n > 2$ are potentials of the first class: BoI and EoI intersect at a certain $\Phi_{\max}$ and the area enclosed is finite. The intersection point is

$$\Phi_{\max} = \frac{1}{4n^2}\left(\frac{3\omega - 2}{\omega}\right)^{n/(n-2)}\left(\frac{32\pi^2}{\lambda n^3}\right)^{1/(n-2)}. \tag{8}$$

Following the same trend as before, the σ field is taken by quantum jumps to the largest values accessible, and we see from Fig. 3 that the quantum scale BoI will wedge in the field σ as it approaches the intersection point, for which the beginning and end of inflation coincide. Therefore, the most typical value of the fields will be that of the intersection point, or strictly speaking, spread over

*The introduction of a non-minimal coupling $\xi\sigma^2 R$ was suggested by [11] to remedy this situation and transform *class II* powerlaw potentials into *class I*, although [11] did not note that $n > 2$ are in fact per se *class I* regardless of non-minimal coupling.

its neighbourhood, as the intersection point as such is a region of measure zero, and the classical path through it has no length. The important prediction of the theories $n > 2$, as all *class I* potentials, is that the crossing of BoI and EoI yields typical values of the fields that are finite.

Class I potentials therefore predict a distribution of M_P at the end of inflation that spans a range of values that is bounded above by $M_P(\Phi_{\max}) < \infty$. This prediction is perfectly consistent with our own observable universe, within a very broad range of parameters (λ,ω,n) allowed by the constraint $M_P(\text{observed}) \lesssim M_P(\Phi_{\max})$. All finite values of M_P satisfying this relation can take place with non-negligible probability, and therefore the scenario is a plausible one. However, *class II* potentials are problematic to reconcile with a finite value of M_P. In these cases, such as $n = 1, 2$, there are still regions that cross EoI at finite values of the fields, though their volume is negligible with respect to other regions for which the fields are arbitrarily large. Such scenarios are inconsistent with the observable universe on the basis of arguments of likelihood and can only be rescued by invoking very strongly the anthropic principle.

ACKNOWLEDGMENTS

The author is grateful to Reza Tavakol, Bernard Carr and Anupam Mazumdar for interesting discussions.

[1] J. Bardeen, P.J. Steinhardt and M.S. Turner, Phys. Rev. D **28**, 679 (1983)

[2] A.S. Goncharov, A.D. Linde and V.F. Mukhanov, Int. J. Mod. Phys. **A2**, 561-591 (1987); A.D. Linde, *Particle Physics and Inflationary Cosmology*, Harwood, Chur, Switzerland (1990); K.A. Olive, Phys. Rep. **190**, 307 (1990)

[3] A.D. Linde, Phys. Lett. B **175**, 395 (1986); Physica Scripta **T15**, 169 (1986); Phys. Today **40(9)**, 61 (1987); A.D. Linde, Nucl. Phys. **B372**, 421 (1992); A.D. Linde, D.A. Linde, and A. Mezhlumian, Phys. Rev. D **49**, 1783 (1994); for a review see [2] and A.R. Liddle and D.H. Lyth, Phys. Rep. **231**, 1 (1993); A.D. Linde, in Proc. of the Erice School "Highlights: 50 Years Later," (1997)

[4] A.A. Starobinsky, in *Current Topics in Field Theory, Quantum Gravity and Strings*, eds. H.J. de Vega and N. Sánchez, Lecture Notes in Physics 206, Springer, Heidelberg (1986)

[5] S.J. Ray, Nucl. Phys. **B284**, 706 (1987); M. Sasaki, Y. Nambu and K. Nakao, Phys. Lett. B **209**, 197 (1988); Nucl. Phys. **B308**, 868 (1988); M. Mijić, Phys. Rev. D **42**, 2469 (1990)

[6] A.D. Linde and A. Mezhlumian, Phys. Lett. B **307**, 25 (1993)

[7] A.D. Linde, D.A. Linde and A. Mezhlumian, Phys. Lett. B **345**, 203 (1995)

[8] T. Singh and T. Singh, Int. J. of Mod. Phys. **A2**, 3, 645 (1987); J.D. Barrow, Phys. Rev. D51, 2729 (1995)

[9] A. Zee, Phys. Rev. Lett. **42**, 417 (1979); B.L. Spokoiny, Phys. Lett. **147B**, 39 (1984); R. Fakir and W.G. Unruh, Phys. Rev. D **41**, 1792 (1990)

[10] D. La and P.J. Steinhardt P.J., Phys. Rev. Lett. **62**, 376 (1989); D. La, P.J. Steinhardt and E. Bertschinger, Phys. Lett. B **231**, 231 (1989); J.D. Barrow and K. Maeda, Nucl. Phys. B **341**, 294 (1990); J. Garcia-Bellido, A.D. Linde and D.A. Linde, Phys. Rev. D **50**, 730 (1994)

[11] J. García–Bellido and A. D. Linde, Phys. Rev. D **52**, 6730 (1995)

[12] M.P. Susperregi, Phys. Rev. D **55**, 560 (1997)

[13] M.P. Susperregi, gr-qc/9712031, "How typical is GR in Brans-Dicke chaotic inflation?"